arXiv:1210.3905v1 [nucl-th] 15 Oct 2012

Heavy-quark transport coefficients in a hot viscous quark-gluon plasma medium

Santosh K. Das [a], Vinod Chandra [b], and Jan-e Alam [c]
[a] *Department of Physics, Yonsei University, Seoul, Korea*
[b] *Istituto Nazionale di Fisica Nucleare (INFN) Sezione di Firenze, Italy and*
[c] *Variable Energy Cyclotron Centre, 1/AF, Bidhan Nagar , Kolkata - 700064*

(Dated: October 16, 2012)

The heavy-quark (HQ) transport coefficients have been estimated for a viscous quark-gluon plasma medium, utilizing a recently proposed quasi-particle description based on realistic QGP equation of state (EoS). Interactions entering through the equation of state significantly suppress the temperature dependence of the drag coefficient of QGP as compared to that of an ideal system of quarks and gluons. Inclusion of shear and bulk viscosities through the corrections to the thermal phase space factors of the bath particles alters the magnitude of the drag coefficient and the enhancement is significant at lower temperatures. The competition between the effects of realistic EoS and dissipative corrections through phase space factor the former eventually dictate how the drag coefficient would behave as a function of temperature, and how much quantitatively digress from the ideal case. The observations suggest significant impact of both the realistic equation of state, and the viscosities, on the HQs transport at RHIC and the LHC collision energies.

PACS: 25.75.-q; 24.85.+p; 05.20.Dd; 12.38.Mh

Keywords: Heavy quark transport; Quasi-particle model; Effective fugacity; Drag and diffusion; Equation of state; Viscous QGP

I. INTRODUCTION

Experimental heavy-ion collision (HIC) program at Relativistic Heavy Ion Collider (RHIC) indicates the production of a liquid like state of the matter where the properties of the system is governed by quarks and gluons. Such a state of matter is called as quark-gluon plasma (QGP). Two of the most striking finding of the RHIC which lead to the imposition of strongly interacting liquid-like behaviour are *viz.*, the large elliptic flow shown by QGP, and the jet quenching phenomenon [1]. Preliminary results from LHC [2–4] confirm the above mentioned observations regarding the QGP. The strongly coupled picture of the QGP is found to be consistent with the lattice simulations of the hot QCD equation of state (EoS) [5, 6], which predict a strongly interacting behavior even at temperatures which are of the order of a $\sim 2T_c$, where T_c is the temperature for the quark-hadron transition. Therefore, it calls for the adaption of a realistic equation of states for the QGP (such as obtained form lattice QCD) in the theoretical and phenomenological investigations regarding its bulk and transport properties. We shall closely follow this point of view in the present analysis in the later sections.

On the other hand, hadrons containing HQs (c, $\bar{c}$ and b, $\bar{b}$) are of great interest in investigating the properties of the QGP, since their physical properties get significantly modified while traveling through the QGP medium. This fact has been reflected in the particle spectra at RHIC, and the LHC energies. Further, HQ thermalization time is larger than gluons and light quarks, and they do not constitute the bulk of the QGP. Since their formation occurs in the early stages of the collisions, they can travel through the thermalized QGP medium, and can retain the information about the interaction with them very effectively. For instance, it is pertinent to ask whether a single $c\bar{c}$ can stay together long enough to form a bound state (say J/ψ) at the hadronization state. To address this, one requires to describe the dynamics of the HQs propagating through the QGP.

Therefore, one can envisage the HQ transport in the QGP medium as follows. The non-equilibrated HQs can travel in the equilibrated QGP medium, and one has to deal the problem within the framework of Langevin dynamics [7]. This is to say that the HQs undertake random motion in the heat bath of the equilibrated QGP. Recall, that the QGP goes through a hydrodynamic evolution process before it reaches the hadronization and subsequently the hadrons freeze-out. The pertinent question to ask is, whether a HQs maintain equilibrium during this entire process of evolution or not. It has been observed [8] within the ambit of Langevin dynamics and pQCD that they do not maintain the equilibrium in the RHIC and the LHC energies.

The most important observable which encode the medium effects carried with them by the HQs while traveling in the QGP, is the nuclear modification factor, R_{AA}. It has been observed that their energy loss in the QGP due to gluon radiation is insufficient to describe the medium modification of the spectrum [9–11]. Therefore, one has to look at the collisions since they have different fluctuation spectrum than radiation, and might contribute significantly as one thought off initially [12]. The collisional effects can be captured well in the HQ drag and diffusion coefficients which have been calculated within weak coupling QCD by several authors. The formalism, and details are offered in Sec. IIA.

The temperature, T and chemical potential, μ_B dependence of the drag and diffusion coefficient enter through the thermal distributions of gluons and light quarks. In the present case, we ignore the μ_B dependence in view of the fact that the QGP produced at RHIC and LHC energies at the mid-rapidity region has negligibly small net baryon density. Therefore, one has to implement the realistic QGP EoS in terms of appropriate form of the thermal distribution functions. Another important aspect is the viscosities (both shear and bulk) of the QGP. One also has to include the viscous modifications in the distribution functions by maintaining the near equilibrium picture so that the mathematical formalism of HQ transport remain intact. This is possible since the QGP posses small shear and bulk viscosities for temperature, $T > T_c$ (T_c is the quark-hadron transition). Both these aspect have been included for the first time while studying the temperature dependence of HQ drag and diffusion coefficients in the QGP medium. Discussion regarding the former is offered in Sec. IIB, and the latter in IIC. Our results suggest that both of these modifications have significant impact on the temperature dependence of HQ transport coefficients. Interestingly, the inclusion of interaction through the EoS modulate the transport coefficients quite significantly as compared to those obtained by employing an ideal EoS.

The paper is organized as follows. In Sec. IIA, formalism of HQ drag and diffusion has been presented. Sec. IIB deals with the quasi-particle description of hot QCD and modeling of thermal distribution functions of gluons, and quarks which constitute hot QGP medium. Sec. IIC deals with the viscous modifications to the thermal distribution functions. In Sec III, results and discussions have been presented for the drag and diffusion coefficients with the realistic EoS. In Sec. IV, this has been investigated including the viscous modifications. Sec. V is devoted to summary and conclusions.

II. HEAVY-QUARK DRAG AND DIFFUSION IN THE VISCOUS QGP

A. Formalism of HQ drag and diffusion

The special role of HQs to characterized QGP stem from the fact that they are produced very early in the collisions and remain extant throughout the evolution and hence can witness the entire evolution of the system. Moreover, HQs do not constitute the bulk part of the system because their mass is considerably larger than the temperature attainable in HIC at RHIC and LHC.

In the present work, we consider the elastic interaction experienced by HQs when propagating in the QGP medium. For the process, $c(p) + l(q) \rightarrow c(p') + l(q')$ (l stands for gluon and light quarks and anti quarks), the drag γ can be calculated by using the following expression [13, 14]:

$$\gamma = p_i A_i / p^2 \qquad (1)$$

where A_i is given by

$$\begin{aligned} A_i = \frac{1}{2E_p} \int \frac{d^3q}{(2\pi)^3 E_q} \int \frac{d^3p'}{(2\pi)^3 E'_p} \int \frac{d^3q'}{(2\pi)^3 E'_q} \\ \frac{1}{g_Q} \sum \overline{|M|^2} (2\pi)^4 \delta^4(p + q - p' - q') \\ f(q)(1 \pm f(q'))[(p - p')_i] \equiv \langle\langle (p - p') \rangle\rangle \end{aligned} \qquad (2)$$

g_Q being the statistical degeneracy of the HQ propagating in the medium. The above expression indicates that the drag coefficient is the measure of the thermal average of the momentum transfer, $p - p'$ due to interaction weighted by the square of the invariant amplitude, $\overline{\mid M \mid^2}$. The factor $f(q)$ denotes the thermal phase space for the particle in the medium. We shall see in the subsequent subsection that $f(q)$ will involve three types of thermal phase space distribution corresponding to the gluons, light-quarks (up and down) and the strange quarks and their anti-quarks. Therefore, $f(q)$ jointly denote these three sectors as,

$$f(q) \equiv \{f_g, f_q, f_s\}, \qquad (3)$$

in the absence of the dissipative effects coming through the shear and the bulk viscosity. It may be recalled here that for zero baryonic chemical potential the thermal phase space for quarks and anti-quarks are same. In the presence of dissipation, we denote it as,

$$f(q) \equiv \{f_{gn}, f_{qq}, f_{ss}\}. \qquad (4)$$

We shall discuss these distributions in the subsequent subsections.

Similar to drag the diffusion coefficient, B_0 can be defined as:

$$B_0 = \frac{1}{4} \left[\langle\langle p'^2 \rangle\rangle - \frac{\langle\langle (p.p')^2 \rangle\rangle}{p^2} \right] \qquad (5)$$

With an appropriate choice of $\mathcal{F}(p')$ both the drag and diffusion co-efficients can be evaluated from the following expression:

$$\begin{aligned} << \mathcal{F}(p) >>= \frac{1}{512\pi^4} \frac{1}{E_Q} \int_0^\infty \frac{q^2 dq d(cos\chi)}{E_q} \hat{f}(q) \\ \frac{w^{1/2}(s, m_Q^2, m_p^2)}{\sqrt{s}} \int_1^{-1} d(cos\theta_{c.m.}) \\ \frac{1}{g_Q} \sum \overline{|M|^2} \int_0^{2\pi} d\phi_{c.m.} \mathcal{F}(p') \end{aligned} \qquad (6)$$

where$s$ is the Mandelstam variable, $E_Q = \sqrt{p^2 + m_Q^2}$ is the energy of the HQ, E_q is the energy of the bath particle $w(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2bc - 2ac$, is the triangular function.

To distinguish among various cases, following notations will be utilized in the subsequent sections. We shall use the notations, γ_η and γ_ζ for the the drag coefficient of the HQs in QGP background with ideal EoS in the presence of shear and the bulk viscosities respectively. On the other hand in the realistic QGP EoS, the same quantities will be denoted as γ^{eos} and γ^{eos} respectively. The ideal counter part (no viscous effects and ideal EoS for QGP) will be denoted as γ_{Id}. Whenever both η and ζ are taken in to account, the drag coefficients will be denoted as $\gamma_{\eta+\zeta}$ in the case of ideal QGP background, and as $\gamma^{eos}_{\eta+\zeta}$ in the case of realistic QGP EoS.

B. The quasi-particle description of hot QCD

Now we discuss the quasi-particle nature of the hot QCD medium which is employed recently in [16]. This description has been developed in the context of the recent (2+1)-lattice QCD EoS [15] at physical quark masses. There are more recent lattice results with the improved actions and refined lattices [6], for which we need to re-look the model with specific set of lattice data specially to define the effective gluonic degrees of freedom. This is beyond the scope of the present analysis. Henceforth, we will stick with the one set of lattice data which is utilized in the model [16].

The model initiates with an ansatz that the Lattice QCD EoS can be interpreted in terms of non-interacting quasi-partons having gluon and quark effective fugacities z_g and z_q respectively which encode all the interaction effects. In this approach, the hot QCD medium is divided into two sectors, *viz.*, the effective gluonic sector, and the matter sectors (light and strange quark). The former refers to the contribution of gluonic action to the pressure which also involves contributions from the internal fermion lines. On the other hand, latter involve interactions among quark, anti-quarks, as well as their interactions with gluons. The ansatz can be translated to the form of the equilibrium distribution functions, $f_{eq} \equiv \{f_g, f_q, f_s\}$ (this notation will be useful later while writing the transport equation in both the sector in compact notations) as follows,

$$\begin{aligned}
f_g &= \frac{z_g \exp(-\beta p)}{\Big(1 - z_g \exp(-\beta p)\Big)}, \\
f_q &= \frac{z_q \exp(-\beta p)}{\Big(1 + z_q \exp(-\beta p)\Big)}, \\
f_s &= \frac{z_q \exp(-\beta\sqrt{p^2+m_s^2})}{\Big(1 + z_q \exp(-\beta\sqrt{p^2+m_s^2})\Big)},
\end{aligned} \tag{7}$$

where $p = |\vec{p}|$, m_s denotes the mass of the strange quark(which we choose to be $0.1GeV$), and $\beta = T^{-1}$ denotes inverse of the temperature. We use the notation $\nu_g = 2(N_c^2 - 1)$ for gluonic degrees of freedom, $\nu_q = 2 \times 2 \times N_c \times 2$ for light quarks, $\nu_s = 2 \times 2 \times N_c \times 1$ for the strange quark for $SU(N_c)$. Here, we are dealing with $SU(3)$, so $N_c = 3$. Since the model is valid in the deconfined phase of QCD (beyond T_c), therefore, the mass of the light quarks can be neglected as compared to the temperature.

Clearly, the determination of f_g, f_q, and f_s require the temperature dependence of the free parameters z_g and z_q respectively. To do this, we rewrite lattice QCD pressure symbolically as,

$$P = P_g + P_{qg}, \tag{8}$$

where P is full (2+1)-flavor lattice QCD pressure, P_g is the contribution coming from the gluonic action alone and P_{qg} is the remaining part after subtracting P_g from P. The fraction P_g, is utilized to define effective gluonic (quasi-gluons) degrees of freedom, and P_{qg} to effective quark-antiquark degrees of freedom through an effective Grand canonical partition function $Z = Z_g \times Z_{qg}$ via the relation $P\beta V = \ln(Z)$ as follows,

$$\begin{aligned}
P_g\beta V &= \frac{V\nu_g}{8\pi^3}\int d^3p \ln(1 - z_g\exp(-\beta p)) \\
P_{qg}\beta V &\equiv (P - P_g)\beta V = \frac{V\nu_q}{8\pi^3}\int d^3p \ln(1 + z_q\exp(-\beta p)) \\
&+ \frac{V\nu_s}{8\pi^3}\int d^3p \ln(1 + z_q\exp(-\beta\sqrt{p^2+m^2})).
\end{aligned} \tag{9}$$

One can read-off Z_g and Z_{qg} which are their standard definitions for bosons and fermions. We know the left hand side of these equation from lattice QCD, and by calculating roots of these equations at each temperature in lattice QCD, we can determine z_g and z_q as a function of temperature. This has been done exactly numerically. Note that we are working at zero baryon chemical potential, therefore quark and antiquark distribution function are the same. This fact has been taken care of by counting the anti-quark degrees of freedom in ν_q and ν_s.

It is worth emphasizing that the effective fugacity is not merely a temperature dependent parameter which encodes the hot QCD medium effects. Its physical significance is reflected through the modified dispersion relation both in the gluonic and matter sector obtained from the thermodynamic relation of energy density $\epsilon = -\partial_\beta \ln(Z)$. One thus find that the effective fugacities modify the single quasi-parton energy as follows,

$$\begin{aligned}
\omega_g &= p + T^2\partial_T ln(z_g) \\
\omega_q &= p + T^2\partial_T ln(z_q) \\
\omega_s &= \sqrt{p^2+m^2} + T^2\partial_T ln(z_q),
\end{aligned} \tag{10}$$

and this lead to the new energy dispersions for gluons (ω_g), light-quark antiquarks (ω_q), and strange quark-antiquarks, (ω_s). These dispersion relations can be explicated as follows. The single quasi-parton energy not only depends upon its momentum but also gets contribution from the collective excitations of the quasi-partons. The second term is like the gap in the energy-spectrum due to the presence of quasi-particle excitations. This makes the model more in the spirit of the Landau's theory of Fermi -liquids. A detailed discussions of the interpretation and physical significance of z_g, and z_q, we refer the reader to [16].

C. Viscous QGP:modification to thermal distributions

Viscosities usually modifies the equilibrium thermal distributions as follows,

$$f = f_{eq} + \delta f_{\eta} + \delta f_{\zeta}. \tag{11}$$

Here, f jointly denote the viscous modified thermal distribution for gluons and quark-antiquarks (in the present case gluons, light quarks and strange quark). Therefore, $f \equiv \{f_{gn}, f_{qq}, f_{qs}\}$. For the viscous corrections δf_{η}, δf_{ζ} we utilize the results of [17] by using only first order terms in the expansion of shear and bulk part of the stress tensor, and choosing the local rest frame of the fluid. Precisely, we have minimally extended the expressions in the present case by substituting the thermal distributions of quarks and gluons as the form obtained in the effective fugacity quasi-particle model. This is solely based on that the quasi-partons in the present case are non-interacting up to their respective effective fugacities that encode all the interactions. This leads to the following expressions,

$$\begin{aligned} f_{gn} &= f_g + \frac{f_g(1+f_g)}{2T^3\tau}\Big[(\frac{p^2}{3} - p_z^2)\frac{\eta}{s} \\ &\quad + \frac{2p^2}{5}\frac{\zeta}{s}\Big] && (12) \\ f_{qq} &= f_q + \frac{f_q(1-f_q)}{2T^3\tau}\Big[(\frac{p^2}{3} - p_z^2)\frac{\eta}{s} \\ &\quad + \frac{2p^2}{5}\frac{\zeta}{s}\Big] && (13) \\ f_{qs} &= f_s + \frac{f_s(1-f_s)}{2T^3\tau}\Big[(\frac{p^2}{3} - p_z^2)\frac{\eta}{s} \\ &\quad + \frac{2p^2}{5}\frac{\zeta}{s}\Big], && (14) \\ & && (15) \end{aligned}$$

where η and ζ are the shear and bulk viscosities of the QGP and s is the entropy density. We will employ the same values of η/s and ζ/s while comparing the effects of ideal and non-ideal EoS with viscosities. For simplicity, we assume same values for η/s and ζ/s for gluons, and quarks. Here, τ is taken to be the thermalization of the QGP.

There are several interesting reports on the estimation of a small value of η/s for QGP, and the emergence of near perfect fluid picture [18–21]. It has been realized in the recent past that bulk viscosity of the QGP is equally significant as the shear viscosity for the temperatures which are close to T_c [22–25]. This is attributed to the large value of the interaction measure, and its behavior with the temperature near T_c [22].

In the present analysis the mass of the strange quarks does not play any prominent role at high temperature ($T \geq T_c$) because $m/T < 1$. In fact, the mass effects are negligibly small (of the order of $(m/T)^2$. Therefore, for the ease in the numerical evaluation the strange quark mass can be ignored.

III. RESULTS AND DISCUSSIONS

Here, we show the temperature dependence of the HQ drag coefficient employing the realistic EOS for the QGP and the dissipative environment mainly induced by the shear and bulk viscosities. Our focus is mainly on the the impact of the the viscous effects, and the realistic QGP EOS. Our aim is served by the ratio in the presence of viscosities and the EOS with respect to that is obtained by assuming the ideal EOS for the QGP medium in which the heavy quarks propagate. These effects are shown in Figs. 1-7.



FIG. 1: γ_{Id} stands for the drag coefficient for ideal case and γ_η stands for drag coefficients when the effect of shear viscosity is included through the phase space of the bath particles. The variation of the ratio of drag coefficients as a function of temperature with different values of η/s at a given thermalization time, τ are shown.

In Fig 1 the effect of shear viscosity on drag coefficient has been depicted as a function of temperature through the ratio between the drag coefficients with shear viscosity to drag coefficient with ideal case. The drag coefficient for zero shear viscosity (ideal case) is higher compared to the case of nonzero shear viscosity in the entire temperature range considered here. Drag coefficient decrease with increase of η/s. The same quantity is displayed in



FIG. 2: Same as Fig. 1 for $\eta/s = 1/(4\pi)$ at two values of τ.



FIG. 3: γ_ζ stands for drag coefficients when the effect of bulk viscosity is included through the phase space of the bath particles. The variation of the ratio of drag coefficients as a function of temperature with different values of ζ/s at a given value of thermalization time τ are depicted.



FIG. 4: Same as Fig. 3 for $\zeta/s = 0.01$ at two different values of thermalization time, τ.



FIG. 5: $\gamma_{\eta+\zeta}$ stands for drag coefficients when the effects of both the shear and bulk viscosities are included through the thermal distribution of the bath particles in evaluating the drag coefficient. The variation of the ratio of the drag coefficients as a function of temperature with different values of ζ/s,η/s and τ are shown.

Fig 2 as a function of temperature for different values of thermalization time at a given value of η/s. It is observed that effect of shear viscosity will decrease if we increase the thermalization time. In Fig 3 the effect of bulk viscosity on drag coefficient has been studied as a function of temperature taking a ratio between the drag coefficients with bulk viscosity to drag coefficient with ideal case. It is found that bulk viscosity increases the drag coefficient. If we increase the ζ/s, drag coefficient is also enhanced as shown in Fig 3. In Fig 4 the effect of thermalization time is displayed. It is found that the effect of bulk viscosity will decrease if we increase the thermalization time. Taking the effect of both the shear and bulk viscosity into consideration the drag coefficient has been calculated ($\gamma_{\eta+\zeta}$) and its variation with the temperature is displayed in Fig 5. As drag coefficients decrease with η/s and increase with ζ/s, it is a interplay between them when both these viscosity effects are included simultaneously. It is found that the low temperature domain is dominated by bulk viscosity where as high temperature part is dominated by shear viscosity and that is why at low temperature drag coefficient is increased and high temperature drag coefficient is reduced as shown in Fig 5. Effect of the EoS on drag coefficient is plotted in Fig 6 by taking a ratio with the ideal case with respect to temperature. It is observed that effect of the EoS suppress the drag coefficient at low temperature. In Fig 7 effects of both shear and bulk viscosities along with the EoS have been presented as a function of temperature at



FIG. 6: γ^{eos} stands for drag coefficients when effect of EoS is considered in evaluating the drag coefficient. The variation of the ratio of the drag coefficients as a function of temperature



FIG. 7: $\gamma^{eos}_{\eta+\zeta}$ stands for drag coefficients when the effects of both the shear and bulk viscosity as well as the effect of EoS are included in evaluating the drag coefficient. The variation of the ratio of the drag coefficients as a function of temperature is depicted here.

given values of τ, η/s, ζ/s. The reduction in the drag is significant at the temperature domain relevant of nuclear collisions at RHIC and LHC energies. Therefore, this results will have crucial effects on the nuclear suppression of flavors measured at these energies.

The effects of phase space corrections due to non-zero shear and bulk viscosities and realistic EoS on diffusion coefficient (B_0) has not been shown explicitly here. However, the effects can be estimated from the drag coefficient (γ) by using the Einstein relation: $B_0 = \gamma m T$.

IV. SUMMARY AND CONCLUSIONS

The drag and diffusion coefficients of the HQs propagating through a thermalized system of quarks and gluons have been evaluated by incorporating the corrections to the phase space factors arising from the non-zero viscosities of the system which restrain the system slightly away from the equilibrium. This corrections seem to be significant for the temperature range expected to be attained in the system produced at HIC at RHIC and LHC collision energies. The effect of realistic EoS on the drag has also been ascertained. The combined effects of viscosities and the EoS reduces the drag significantly. These results will have crucial consequences on the heavy ion phenomenology at RHIC and LHC energies.

Acknowledgements

SKD thanks Payal Mohanty for useful discussions. VC sincerely acknowledges the financial support from INFN, Italy for awarding an INFN postdoctoral fellowship at the Firenze section.

---

[1] J. Adams *et al.* (STAR collaboration), Nucl. Phys. **A757**, 102 (2005); K. Adcox *et al.* (PHENIX Collaboration), *ibid.* **A757**, 184 (2005); Back *et al.* (PHOBOS Collaboration), *ibid.* **A757**, 28 (2005); A. Arsence *et al.* (BRAHMS Collaboration), *ibid.* **A757**, 1 (2005).

[2] M. Krzewicki (ALICE Collaboration), QM-2011, `arXiv:1107.0080v1 [nucl-ex]`.

[3] K. Aamodt *et al.* (The Alice Collaboration), `arXiv:1011.3914 [nucl-ex]`; Phys. Rev. Lett. **105**, 252301 (2010); Phys. Rev. Lett. **106**, 032301 (2011).

[4] T. Hirano, P. Huovinen, Y. Nara, `arXiv:1012.3955[nucl-th]`; X. -F. Chen, T. Hirano, E. Wang, X. -N. Wang, H. Zhang, `arXiv:1102.5614[nucl-th]`.

[5] G. Boyd *et. al*, Phys. Rev. Lett. **75**, 4169 (1995); Nucl. Phys. **B 469**, 419 (1996); M. Panero, Phys. Rev. Lett. **103**, 232001 (2009); F. Karsch, E. Laermann, A. Peikert, Phys. Lett. **B 478**, 447 (2000); M. Cheng *et. al*, Phys. Rev. **D 77**, 014511 (2008); A. Bazavov *et. al*, Phys. Rev. **D 80**, 014504 (2009); M. Cheng *et. al*, Phys. Rev. **D 81**,054504 (2010).

[6] S. Borsanyi *et. al*, JHEP **1009**,073 (2010); JHEP **11**, 077 (2010); Y. Aoki *et al.*, JHEP **0601**, 089 (2006); JHEP **0906**, 088 (2009).

[7] E. M. Lifshitz, and L. P. Pita Pitaevskii, Physics kinetics (Butterworth-Hienemann, Oxford 1981); R. Balescu, Equilibrium and Non-equilibrium Statistical Mechanics (Wiley, New York 1975).

[8] S. K. Das, Jan-e Alam, and Payal Mohanty, Phys. Rev. **C 80**, 054916 (2009).

[9] G. D. Moore, and D. Teaney, Phys. Rev. **C 71**, 064904 (2005).

[10] R. Baier, Y. L. Dokshitzer, A. H., Mueller, and D. Schiff, JHEP **0109**, 033 (2001).

[11] S. Jeon, and G. D. Moore, arXiv:hep-ph/0309332.

[12] M. G. Mustafa, and M. H. Thoma, arXiv:hep-ph/0309332.

[13] B. Svetitsky, Phys. Rev. D **37**, 2484(1988).

[14] G. D. Moore and D. Teaney, Phys. Rev. C **71**, 064904 (2005); H. van Hees, R. Rapp, Phys. Rev. C,**71**, 034907 (2005); S. K Das, J. Alam and P. Mohanty, Phys. Rev. C **82**, 014908 (2010); S. K Das, J. Alam, P. Mohanty and B. Sinha, Phys. Rev. C **81**, 044912 (2010); S. Majumdar, T. Bhattacharyya, J. Alam and S. K. Das, Phys. Rev. C **84** , 044901 (2012); P. B. Gossiaux and J. Aichelin, Phys. Rev. C **78**, 014904 (2008); J. Uphoff, O. Fochler, Z. Xu and C. Greiner, Phys. Rev. C **84** 024908 (2011); W. M. Alberico *et al.*, Eur.Phys.J.C **71** 1666 (2011); Y. Akamatsu, T. Hatsuda and T. Hirano, Phys. Rev. C **79**, 054907 (2009); H. van Hees, M. Mannarelli, V. Greco and R. Rapp, Phys. Rev. Lett. **100**, 192301 (2008), M. G. Mustafa, D. Pal and D. K. Srivastava, Phys. Rev. C **57**, 889 (1998).

[15] M. Cheng *et. al*, Phys. Rev. **D 77**, 014511 (2008).

[16] V. Chandra, V. Ravishankar, Phys. Rev. **D 84**, 074013 (2011).

[17] K. Dusling, and T, Schäfer, Phys. Rev. **C 77**, 034905 (2008); K. Dusling, and D. Teaney, Phys. Rev. **C 85**, 044909 (2012).

[18] H. B. Meyer, Phys. Rev. **D 76**, 10171 (2007); Lacey *et. al*, Phys. Rev. Lett. **98**, 092301 (2007); Zhe Xu and C. Greiner, Phys. Rev. Lett. **100**, 172301 (2008); Zhe Xu, C. Greiner, H. Stoecker, Phys. Rev. Lett. **101**, 082302 (2008); Adare *et. al*, Phys. Rev. Lett. **98**, 172301 (2007); S. Gavin and M. Abdel-Aziz, Phys. Rev. Lett. **97**, 162302 (2006); A. Buchel, Phys. Lett. **B 663**, 286 (2008); P. Kovtun, D. T. Son, A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005).

[19] M. Asakawa, S. A. Bass, B. Müller, Phys. Rev. Lett. **96**, 252301 (2006); Prog. Theor. Phys. **116**, 725 (2007).

[20] V. Chandra and V. Ravishankar, Euro. Phys. J C **59**, 705 (2009).

[21] V. Chandra and V. Ravishankar, Euro. Phys. J C **64**, 63 (2009).

[22] F. Karsch, D. Kharzeev, K. Tuchin Phys.Lett. **B 663**, 217 (2008) ; D. Kharzeev, K. Tuchin, JHEP **09**, 093 (2008).

[23] A. Buchel, Phys. Lett. **B 663**, 286 (2008).

[24] V. Chandra, Phys. Rev. **D 84**, 094025 (2011).

[25] V. Chandra, `arXiv:1202:1255 [nucl-th]`.